

-continued

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 110

cttggttatg ttggcactga tc

22

<210> SEQ ID NO 111

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 111

tgtgttaggg gactggtgga ca

22

<210> SEQ ID NO 112

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 112

catcacccac ttacccccaa aa

22

<210> SEQ ID NO 113

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 113

ataaccgagt cgttctgcca at

22

<210> SEQ ID NO 114

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide

<400> SEQUENCE: 114

tttcagagca ttggccatag aa

22

What is claimed is:

1. A method for treating or preventing cancer in a subject in need thereof comprising administering to the subject an amount of nicotinamide mononucleotide (NMN), or a salt thereof, or a prodrug thereof, effective to increase the level of nicotinamide adenine dinucleotide (NAD⁺) in the subject.

2. The method of claim 1, wherein the NMN or salt thereof, or prodrug thereof, is administered at a dose of between 0.5-5 grams per day.

3. The method of claim 1, wherein the subject is a human.

4. The method of claim 1, wherein the NMN or salt thereof, or prodrug thereof, is administered as a pharmaceutical composition comprising a pharmaceutically acceptable carrier and the NMN or salt thereof, or prodrug thereof.

5. The method of claim 4, wherein the pharmaceutical composition is adapted for oral administration.

6. The method of claim 1, wherein the NMN or salt thereof, or prodrug thereof, is administered orally.

7. The method of claim 2, wherein the NMN or salt thereof, or prodrug thereof, is administered orally.

8. The method of claim 4, wherein the NMN or salt thereof, or prodrug thereof, is administered orally.

9. The method of claim 2, wherein the subject is a human.

10. The method of claim 4, wherein the subject is a human.

11. The method of claim 5, wherein the subject is a human.

12. The method of claim 2, wherein the NMN or salt thereof, or prodrug thereof, is administered as a pharmaceutical composition comprising a pharmaceutically acceptable carrier and the NMN or salt thereof, or prodrug thereof.

13. The method of claim 12, wherein the pharmaceutical composition is adapted for oral administration.

14. The method of claim 12, wherein the NMN or salt thereof, or prodrug thereof, is administered orally.

15. A method for treating or preventing cancer in a subject in need thereof, comprising administering to the subject an amount of NMN, or a salt thereof, effective to increase the level of NAD⁺ in the subject.